

35% Carbamide Peroxide gel의 적용시간에 따른 법랑질 변화

1

, 1

1. 서론

10% carbamide peroxide가
22% 35% car-
bamide peroxide²⁾,
가

10% 15% carbamide peroxide 2
15%
3) 10% 20% carbamide peroxide
2 20%

carbamide peroxide
가 가 가 carbamide peroxide
30~35% hydrogen peroxide
35% carbamide peroxide
가
가 carbamide peroxide 가
가

Heywood Heymann¹⁾ 10% carbamide peroxide 15% carbamide
peroxide
가 35% carbamide peroxide

연락처: 이해진 우) 612-715 부산시 해운대구 반송2동 640-5번지 동부산대학 치위생과
전화: 051-540-3877, 휴대폰: 016-467-6679 E-Mail: onlyhelena@hanmail.net

가 6-

¹⁰⁾. Carbamide peroxide hydro-
gen peroxide urea , 10% car-
bamide peroxide 3.62% hydrogen per-
oxide , 22% carbamide peroxide
7.92% hydrogen peroxide
35% carbamide peroxide
11.7% hydrogen peroxide
6% hydrogen peroxide
⁷⁾
가 ⁸⁾.
carbamide peroxide
가

2.1.1 시편제작

0.1% (Sigma, U.S.A.)
1×1cm 가
24 (Automatic
Polisher; Labopol-1, Struers, Denmark)
800, 1000, 1200,
1500 grit abrasive paper
, 1μm
30

2.2 연구방법

2.2.1 시편변색 및 전처리

가 가 (; pH 2.26±0.02) 24
35% carbamide 가
peroxide 가
가 24 가 , 24 3
4
(Table 1). WHITE

2. 연구재료 및 방법

2.1 연구재료

smile 35% Carbamide Peroxide gel
(WHITE smile GmbH. Inc. Germany)
1 7
1

표 1. The classification bleaching group according to bleaching time

Group	Classification	Application period
1	Control group	Distilled water(D.W.)
2	Experimental group	35% Carbamide Peroxide gel 1hour/day
3	"	35% Carbamide Peroxide gel 2hour/day
4	"	35% Carbamide Peroxide gel 3hour/day

2 , 3

7

bleaching gel

D.W(Distilled water) 가

1

Japan)

3

, 3 , 7

20kV 가

1 μ m

222. 법랑질 표면의 미세경도 측정

Vickers diamond indenter가

(MVK-H100, Hardness Testing

Machine, Akashi Corporation, Japan)

vickers hardness number(VHN)

200gm

10

400

300~400 VHN

3

223 법랑질 표면의 미세구조 분석

(Field emis-

sion scanning electron microscope, Hitachi, Japan)

3

1

2

(Ion sputter, Hitachi,

Japan)

70nm

20kV

224 치아 표면 무기질의 변화

(EDS, S-4300 & EDX-350, Hitachi,

Japan)

1 , 3 , 7

200 μ m

225 자료분석

SPSS

13.0 package program

paired t-test

Duncan

3. 연구성적

3.1 35% Carbamide Peroxide gel의 노출시간에 따른 법랑질의 미세경도 변화

(VHN) 7

1

($p>0.05$).

2 , 3

가

3

($p<0.05$)<Table 2>.


표 2. VHN values of enamel during the process

Group	N	Baseline (24hours staining.)		Treatment days			p-value
				1 day	3days	7days	
D.W	6	233.17 ±3.24	^a	231.12 ±4.35 ^a	228.32 ±8.59 ^a	228.44 ±5.21 ^a	p>0.05
CP 1 hours	6	236.12. ±6.21	^a	235.11 ±5.24 ^a	228.33 ±7.96 ^a	226.58 ±9.21 ^a	p>0.05
CP 2 hours	6	233.78 ±3.69	^a	231.78 ±3.69 ^a	221.58 ±4.59 ^b	212.59 ±10.58 ^c	p<0.05
CP 3 hours	6	235.74 ±4.12	^a	229.74 ±4.12 ^a	210.29 ±12.39 ^c	191.44 ±14.55 ^d	p<0.05

Values are reported as the Mean ± Standard deviation
^{a,b,c,d} The same letter indicates no significant difference at α= 0.05 by Duncan's studentized range test
Statistical comparison by ANOVA test

3.2 35% Carbamide Peroxide gel의 노출시간에 따른 법랑질 표면의 미세 구조 변화

Hitachi, Japan)



<Figure 1>.
Figure 1. SEM images of enamel surfaces. A: 35% Carbamide Peroxide gel 1 hour for 7days, B: 35% Carbamide Peroxide gel 2 hours for 7days, C: 35% Carbamide Peroxide gel 3 hours for 7days, D: Sound tooth (baseline), E: Enamel with cola staining (baseline)

(Field emis-
sion scanning electron microscope,

3.3 35% Carbamide Peroxide gel의 노출시간에 따른 법랑질 표면의 무기 질 변화

표 3. Contents of calcium and phosphate in enamel surface for 7 days

Group	N	Baseline (24hours staining)				7 days			
		Ca		P		Ca		P	
D.W	6	33.46±1.25	^a	17.54±0.25	^b	32.56±1.22	^a	17.12±0.46	^b
CP 1 hours	6	34.18±2.58	^a	18.09±1.02	^b	32.05±1.46	^a	18.02±0.57	^b
CP 2 hours	6	31.45±3.97	^a	18.24±0.55	^b	30.86±1.25	^a	17.46±1.33	^b
CP 3 hours	6	34.05±2.56	^a	17.58±0.48	^b	27.27±1.99	^a	14.39±0.28	^b

Values are reported as the Mean ± Standard deviation
^{a,b} The same letter indicates no significant difference at α= 0.05 by Duncan's studentized range test

Ca, P

¹²⁾

3

가

35% Carbamide Peroxide

($P>0.05$)<Table 3>.

4. 중괄 및 고안

1

가

가

가

1

2

3

Ol tu Gurgan⁶⁾ 10%, 16% Carbamide Peroxide

가

35% Carbamide

Peroxide

6-10)

가

35%

가

Carbamide Peroxide

1

(SEM)

11).

가

Ben-Awar ¹³⁾, McGuckin

¹⁴⁾

Nathoo

¹⁵⁾

Ca, P

16)

3.

Ca, P

35% Carbamide Peroxide gel

($P>0.05$).

35% carbamide peroxide(CP)

가

in vitro

가

가

in vivo, in situ

가

참고문헌

- 가
5. 결론
- 가
- 35% Carbamide Peroxide
- 가
- 1.
- 7
- 가
- 2
- ($P<0.05$).
- 2.
1. Haywood VB, Heymann HO. Nightguard vital bleaching. Quintessence Int 1989; 20(3):173-176.
2. Matis BA. Degradation of gel in tray whitening. Compend Cotin Educ Dent Suppl 2000;28:28, 31-35.
3. Kihn PW, Barnes DM, Romberg E, Peterson K. A clinical evaluation of 10 percent vs 15 percent carbamide peroxide tooth whitening agents. J Am Dent Assoc 2000;131(10):1478-1487.
4. Jones AH, Diaz-Arnold AM, Vargas MA, Cobb DS. Colorimetric assessment of laser and home bleaching techniques. Esthet Dent 1999;11(2):87-94.
5. Leonard RH, Sharma A, Haywood VB. Use of different concentrations of carbamide peroxide for bleaching teeth: an in vitro study. Quintessence

- Int 1998;29(8):503-507.
6. Oltu U, Gurgan S. Effects of three concentrations of carbamide peroxide on the structure of enamel. *J Oral Rehabil* 2000;27(4):332-340.
 7. Lewinstein I, Hirschfeld Z, Stabholz A, Rotstein I. Effect of hydrogen peroxide and sodium perborate on the microhardness of human enamel and dentin. *J Endod* 1994;20(2):61-63.
 8. Heling I, Parson A, Rotstein I. Effect of bleaching agents on dentin permeability to *Streptococcus faecalis*. *J Endod* 1995;21 (11):540-542.
 9. Rotstein I, Dogan H, Avtron Y, Shemesh H, Stein berg D. Protective effect of copalite surface coating on mercury release from dental amalgam following treatment with carbamide peroxide. *Endod Dent Traumatol* 2000;16(3):107-110.
 10. Bitter NC, Sanders J. Electron microscopy study of four bleaching agents on enamel surfaces *J Dent Res* 1992;71(4):600-602.
 11. Joiner A, Jones NM, Raven SJ. Investigation of factors influencing stain formation utilizing an in situ model. *Adv Dental Res* 1995;9:471-476.
 12. Zalkind M, Arwaz JR, Goldman A, Rotstein I. Surface morphology changes in human enamel, dentin and cementum following bleaching. *Endod Dent Traumatol* 1996;12:82-88.
 13. Ben-Amar A, Liberman R, Gorfil C, Bernstein Y. Effect of mouthguard bleaching on enamel surface. *J Am Dent* 1995 ;8(1):29-32.
 14. McGuckin RS, Babin JF, Meyer BJ. Alterations in human enamel surface morphology following vital bleaching. *J Prosthet Dent* 1992;68(2):754-760.
 15. Nathoo SA, Chmielewski MB, Kirkup RE. Effect of Colgate Platinum Professional Tooth whitening System on microhardness of enamel, dentin, and composite resins. *Compendium. Suppl* 1994;17:627-630.
 16. Kown YH, Son KH, Kim GC, Kim HI. Effect of 35% Carbamide Peroxide on Enamel: Hardness, Structure and Composition. *J Korean Res Soc Dent Mater* 2003;30(4):443-449.

Abstract

Surface change of enamel according to application time of 35% Carbamide Peroxide

Hye-Jin Lee, Min-Young Kim

Dept of Dental Hygiene, Dongpusan College University

Key words: Enamel surface, 35% Carbamide Peroxide, Microhardness

The purposes of this study were to examine the effect of 35% Carbamide Peroxide(CP) bleaching agent on the changes in physical and chemical characteristics of tooth. The effect of bleaching agent on enamel was analyzed using Hardness test, SEM and EDS. The microhardness between bleached groups after bleaching showed statistically significant difference according to the paired t-test. The bleached enamel surface showed apparent morphological changes compared to the enamel, which was stored in distilled water only. The difference of the total mineral contents for the distilled water and Carbamide Peroxide did not show statistical significance.

These results demonstrated that bleaching using 35% Carbamide Peroxide were adversely affects application time of experimental group and may confirm the safety of using these agents for a short time in dentist-monitored bleaching.

- 2008. 9. 30

- 2008. 11. 24

- 2008. 12. 15