Journal of Korean Society of Dental Hygiene

Original Article

The correlation between physique and dental arch size

Seung-Hun Lee[®]• In-Suk Park[®]

Department of Dental Hygiene, Cheongam College

Corresponding Author: Seung-Hun Lee, Department of Dental Hygiene, Cheongam College, 1641, Noksae-or, Suncheon, 577997, Korea. Tel: +82-61-740-7382, Fax: +82-61-740-7418, E-mail: smili@hanmail.net



Received: November 12, 2019 Revised: January 10, 2020 Accepted: January 13, 2020

ABSTRACT

Objectives: We analyzed the correlation between physique and size of the dental arch of college students with normal occlusion. **Methods:** Ninety-nine college students filled out the prepared questionnaire. The length and width of the dental arch of the students was measured using a plaster model after taking an impression. The data were analyzed using the t-test and correlational analysis. **Results:** The maxilla arch length was 3.70 mm longer and the inter-molar width of the maxilla was 3.06 mm longer in men (both *p*<0.01) than in women. Additionally, the mandible was 3.66 mm longer in men as compared to those in the women (*p*<0.01). As height increased, there was increase in the body weight (α =0.683, *p*<0.01), maxilla arch length (α =0.373, *p*<0.01), molar width of the maxilla (α =0.214, *p*<0.05), and the mandible (α =0.280, *p*<0.01). The greater the weight increase, the greater the maxillary arch length increase (α =0.392, *p*<0.01), and the greater the molar width of the maxilla (α =0.336, *p*<0.01) and mandible (α =0.502, *p*<0.01) increase. **Conclusions:** As physical size (both height and weight) increased, the maxillary length and molar width also increased. Based on the results of this study, the cause of malocclusion, form and size of the dental arch, and stable occlusion can be used as basic data.

Key Words: Body height, Body weight, Dental arch, Somatotype, Student

Introduction

The dental arch is an interesting area of study in dentistry because of the variety in its form. This is especially important because of the basic data on the treatment plane of the tooth correction and the orthognathic surgery [1]. It has been studied about the size, position of tooth and dental arch to make a stable, functional, and aesthetic dental arch [2-4]. There are studies on the modality of malocclusion correction according to regional differences in anatomy [2], the effect of tooth size and dental arch size on the crowding [3], and comparison of arch forms between Korean and Japanese patients with malocclusion [4].

Since the 1970s, rapid development has improved diet, resulting in a change in the physique, body shape, and facial shape of individuals [5]. In a study by Kim [6], the average height of a 14-year-old was 152.0 cm in males and 151.0 cm in females in 1970, and 162.5 cm in males and 156.6 cm in females in 1990. Body weight was 43.7 kg in males and 43.8 kg in females in 1970, and 51.9 kg in males and 49.6 kg in females in 1990. According to a recent study [7], the average height and weight of Korean children and adolescents between 2006 and 2015 were 167.9 cm in males and 159.2 cm in females, and 58.4 kg and 52.8 kg in males and females, respectively. Height and weight are constantly increasing.

Angle [8] stated that teeth in normal occlusion are located on a line matching the shape and position of the arch, and Sved [9] stated that the dental arch was shaped by the function of each tooth, and the teeth were accordingly arranged on the spherical surface. The size discrepancy of the jaw and teeth can affect the dental arch and lead to teeth crowding and malocclusion [10]. Orthodontic treatment improves teeth arrangement and occlusion, but post-treatment stability is affected by the size of the dental arch [11,12].

Data on growth and changes in the dental arches are important for analyzing the physique changes with age and understanding how these changes are related to the dental arches. As the physique improves, size of the dental arches is predicted to change. Therefore, it is necessary to analyze the size of the dental arch according to the physique and use the results in the treatment plan.

Recently, various studies on the dental arch have been reported [1,4,11,13-19]. There are studies on the size of the dental arch according to malocclusion [4,13], size of the dental arch according to race [4,14], changes in the dental arch after orthognathic surgery [1,15], effects of tooth extraction on the dental arch [11,16], changes in the dental arch according to age [17,18], etc. However, studies on the size of the dental arch according to physique are insufficient. The previous study was limited to females [19]. Therefore, we aimed to measure the physique of males also and analyze the correlation between the dental arch size and physique.

Methods

1. Study subjects

The study subjects were 120 students of the C college in Suncheon city. The purpose and method of the study was explained, and they were asked to fill out the questionnaire, to which signed the informed consent. The survey period was five days, from November 26 to 30, 2018, and the questionnaires were collected directly by the researcher. Ninety-nine questionnaires were used for the final data analysis, excepting those of 17 experienced patients undergoing orthodontic treatment with tooth extraction, and those of four missing and incomplete responses. The plaster models of their dental arch were made after impression taking.

The minimum number of study samples was calculated as 84 using the G^{*}power 3.1[20,21] program by correlation analysis with a significance level of 0.05, test power of 80%, and 0.3 effect size. Assuming

the rate of elimination as 20%, 120 subjects were suitable for the analysis. The size effect was referenced from the results of the study by Han et al. [19], which analyzed the correlation between size of the dental arch and physique of the dental students.

This study was approved by the 00 Institutional Review Board (CA17-190326-HR-001-01).

2. Study method

The questionnaire consisted of eight questions, including four questions on general characteristics including weight and height, respiration, orthodontic treatment experience, tooth extraction experience, and favorite food. The items were modified and adapted according to the study purpose by referring to the survey questionnaire by Han et al. [19].

The surveyor made plaster models after recording impressions of the dental arches of the subjects with alginate impression material. The measurements are shown in <Fig. 1>. Length of the dental arch is the distance from the incisal edge of the central incisor to the distal surface of the first molar; intercanine width is the distance between tips of cusps of the left and right canines; inter-molar width is the distance between the mesiobuccal cusps of the left and right first molars. Size of the dental arch was measured using a vernier caliper (Digimatic caliper, Sincon, China).



Fig. 1. Measurements of the dental arch 1: arch length, 2: inter-canine width, 3: inter-molar width

3. Data analysis

Independent t-test was used to compare the differences between the length and width of the dental arches according to the general characteristics. Pearson correlation coefficient was calculated for the correlation between the physique and size of the dental arch. The statistical significance level was p-value of less than 0.05. The data were analyzed using SPSS (ver. 18.0, Chicago, Illinois, USA).

Results

1. General characteristics of the study subjects

The study subjects were 99 students of the Cheongam College, and their details are shown in Table 1. Their average age was 19.2 years, and average weight and height were 66.99 kg and 169.69 cm, respectively. Of the subjects, 83.8% had a nasal breathing and 62.6% preferred solid food.

(11 00)

Table 1. Characteristics of the study subjects									
Characteristics	Division	Mean SD Characteristics		Division	Ν	%			
Age		19.2	0.91	Gender	Male	50	50.5		
Weight	Male	75.16	9.82		Female	49	49.5		
	Female	58.65	4.91	Respiration	Oral	16	16.2		
	Total	66.99	11.35		Nasal	83	83.8		
Height	Male	177.42	5.16	Favorite food	Solid food	62	62.6		
	Female	161.80	2.60		Soft food	37	37.4		
	Total	169.69	8.85						

Table 1 Characteristics of the study subjects

2. The length and the width of dental arches according to the general characteristics

The length and width of the dental arches according to the general characteristics have been presented in Table 2. Males had 3.70 mm longer maxillary arch than females (p < 0.01), 3.06 mm longer inter-molar distance (p<0.01) in the maxilla, and 3.66 mm longer inter-molar distance in the mandible (p < 0.01). Other variables did not show a statistically significant difference between the sexes.

The subjects whose favorite was soft food had 2.07 mm longer mandibular arch length than the subjects whose favorite was solid food (p < 0.05), and 1.06 mm longer inter-canine distance in the mandible (p < 0.05). There were no statistically significant differences in other variables according to the favorite food.

3. Correlation between the physique and size of dental arch

Correlation between the physique and size of the dental arch is shown in Table 3. As height increases, the weight also increases (α =0.683, p<0.01). In addition, the maxillary arch length (α =0.373, p<0.01) and the molar width of the maxilla (α =0.214, p<0.05) and mandible (α =0.280, p<0.01) also increases. More the increase in weight, more the maxillary arch length increase (α =0.392, p<0.01). Molar width of maxilla (α =0.336, p<0.01) and mandible (α =0.502, p<0.01) also showed an increase.

Variables a	Maxilla arch length	p^{*}	Mandible arch length	<i>p</i> *	Maxilla inter- canine width	p^{*}	Mandible inter- canine width	p^{*}	Maxilla inter-molar width	p^{*}	Mandible inter-molar width	p^{*}
Gender												
Male 4	43.08±5.33	< 0.001	37.40±3.72	0.792	40.80 ± 3.10	0.825	33.30 ± 1.74	0.573	60.46 ± 3.71	< 0.001	56.95±3.34	< 0.001
Female 3	39.38±2.11		37.20±3.65		40.66±3.11		33.02±2.97		57.40±2.02		53.30±2.06	
Favorite food	d											
Solid food	41.10±4.35	0.667	36.51±3.15	0.010	40.89±3.42	0.514	32.75±2.52	0.034	58.72±3.56	0.399	54.97±3.28	0.524
Soft food	41.50±4.66		38.57±4.11		40.47±2.49		33.81±2.13		59.31±3.36		55.41±3.41	
Total 4	41.25±4.45		37.30±3.67		40.73±3.09		33.16±2.42		58.95±3.35		55.14±3.32	

Table 2. Length and the width of dental arch according to the general characteristics

Unit: Mean±SD

* by the independent t-test at α <0.05

Table 3. Correlation between the physique and the dental arch

Variables	Weight	Height	Maxilla arch length	Mandible arch length	Maxilla inter- canine width	Mandible inter- canine width	Maxilla inter- molar width	Mandible inter- molar width
Weight	1							
Height	0.683**	1						
Maxilla arch length	0.373**	0.392**	1					
Mandible arch length	0.106	0.105	0.461**	1				
Maxilla inter-canine width	-0.081	-0.017	0.235*	0.255*	1			
Mandible inter-canine width	0.171	-0.024	0.071	0.285**	0.315**	1		
Maxilla inter-molar width	0.214*	0.336**	0.197	0.079	0.398**	0.271**	1	
Mandible inter-molar width	0.280**	0.502**	0.221*	0.157	0.220*	0.249*	0.853**	1

** p<0.01 by Pearson's correlation analysis

Discussion

Rapid growth of the economy and improved diet have led to a significant increase in height and weight when compared to the past in Korea. As physique increases, size of the dental arch will change too. As the physique grows, the dental arches grow, and the mismatch of the maxillary and mandibular teeth causes malocclusion. Growth of the body and size of the dental arch are important because they are the basis for diagnosing and treating malocclusion.

The average age of the subjects included in the study was 19.2 years, height was 169.7 cm (male 177.4 cm, female 161.8 cm), and weight was 66.9 kg (male 75.2 kg, female 58.7 kg). In 2015, a 19-year-old Korean had an average height of 168.4 cm (male 173.4 cm, female 160.7 cm) and weight of 66.1 kg (male 71.1 kg, female 58.3 kg) [20]. In the study by Han et al. [19], females had a height of 163.2 cm and weight of 52.3 kg. The height of females in this study was lower than that in the study of 2013 and somewhat higher than the average height in Korea in 2015. The weight was also more. Although there are some differences, age is not considered a significant variable because all subjects in the study were adults.

In the mandible of females, length of the arch was 37.20 mm, inter-canine width was 33.02 mm, and distance between the molars was 53.30 mm. In the study by Han et al. [19], these were 31.8 mm, 26.8 mm, and 40.5 mm, respectively. These values are respectively 23.7 mm [21], 26.6 mm, 45.8 mm in the study by Hwang et al.[21]. The size of the arch form was more than that of the previous study [20,21].

Similar to the previous studies [20,21], the width was larger than the length. Hwang et al. [21] reported that the width of the dental arch narrowed due to changes in eating habits, and predicted that this trend will progress in future. These changes in the arch form need to be further investigated.

Variations in the size of the dental arch by sex showed greater size in males than in females. However, there was a statistically significant difference in the maxillary arch length (p<0.01) and the maxillary molar width (p<0.01). These results may be justified by the fact that physique of males is greater than that of females. Some studies have reported no differences in the size of dental arches between sexes [22-24], while others have shown that males have larger arch length [25,26]. Follow-up studies on various age groups are needed to verify the differences in the size of the dental arches by sex.

No large difference in the size of the arch according to the favorite food was found. There was a statistically significant difference in the arch length of the mandible and mandibular inter-canine width, but the difference was small. There was no statistically significant difference in the previous study [19]. Processed foods have reduced the demand for masticatory muscles and teeth. In the survey of medieval skulls, modern humans were found to have smaller depth and width of dental arches than their ancestors [27,28]. Follow-up studies are needed to determine the size of the dental arch according to the favorite food during the growing period.

As the physique increased, size of the dental arch increased. In particular, there was a statistically significant difference in the length of the arch and molar width. In the study by Han et al. [19], there was a statistically significant correlation between the weight and size of the dental arch. As the weight increased, inter-canine and inter-molar widths increased. However, there was no correlation with height. There was no correlation between height and weight in the previous study; however, in this study, weight increased as height increased.

This study has a limitation of generalization, as measurements were performed only for some college students, and the physique of males and females was not similar. However, unlike previous studies [19], the males were also measured, except those who had undergone extraction and orthodontic treatment that could affect the size of the arch. Follow-up studies will need to investigate different age groups, breathing methods, and malocclusion types. Based on the results of this study, the cause of malocclusion, form, and size of the dental arch, stability of occlusion, and oral health can be used as basic data.

Conclusions

In this thesis, we analyzed the correlation between college students' physique and dental arch size. Ninety-nine subjects completed the questionnaire and underwent impression taking. The t-test and correlation analysis were used for analysis.

The measurements and analysis results are as follows.

1. In males, the maxillary arch length (p<0.01) and inter-molar width of the maxilla and mandible (p<0.01) were higher than that of females.

2. Differences in the dental arch size according to favorite food was not significant.

3. As height increased, the maxillary arch length increased (α =0.373, p<0.01), and molar width of the

maxilla (α =0.214, p<0.05) and the mandible (α =0.280, p<0.01) also increased.

4. More the weight increase, more the increase in maxillary arch length (α =0.392, p<0.01) and molar width of the maxilla (α =0.336, p<0.01) and mandible (α =0.502, p<0.01).

The size of the dental arch in males is larger than that in females. As the physique increases, maxillary length and molar width increases. These results can be used as basic data to study the cause of malocclusion, form, and size of the dental arches, stability of occlusion, and oral health.

Conflict of interest

The authors declared no conflicts of interest.

Authorship

Conceptualization: SH Lee; Data collection: IS Park; Formal analysis: SH Lee; Writing – original draft: SH Lee; Writing – review & editing: SH Lee, IS Park

References

- [1] Ryu KS, Lee BS, Kim YG, Kwon YD, Choi JC, Ohe JY. Surgical and orthodontic treatment of skeletal class III featuring severe transversal and sagittal discrepancy : A case report. J Korean Assoc Maxillofac Plast Reconstr Surg 2013;35(2):124-9. https://doi.org/10.14402/ jkamprs.2013.35.2.124
- [2] Sohn BW, Hwang CJ, Hwang HS. Modality of malocclusion according to regional difference using hld(handicapping labiolingual deviation), tpi(treatment priority index) and hmar(handicapped malocclusion assessing record). Korean J Orthod 1993;23(1):17-35.
- [3] Ahn BK. A statistical study on the effect of tooth size and dental arch size upon the crowding. Korean J Orthod 1990; 20(2):217-26.
- [4] Lee CH, Mo SS, Kang YG, Nojima K, Kim YH, Kook YA. Comparison of arch forms between Korean and Japanese in Class 1, II, and III malocclusion. Korean J Orthod 2007;37(5):364-75.
- [5] Leem HS, Lee KJ. Facial measurements and analysis of middle school students in Seoul. Korean J Vis Sci 2003;5(2):63-72.
- [6] Kim JS. Perspective on population characteristics and health problems of Koreans in 21st century. Korean J Preventive Medicine 1994;27(2):175-85.
- [7] Kwon EJ, Nah EH. Secular trends in height, weight and obesity among Korean children and adolescents in 2006-2015. Korean J Health Educ Promot 2016;33(2):1-13. https:// doi.org/10.14367/kjhep.2016.33.2.1
- [8] Angle EH. Treatment of malocclusion of the teeth. 7th ed. Philadelphia: The S.S. White Dental Manufacturing Co; 1907.
- [9] Sved A. The application of engineering methods to orthodontics. Am J Orthod 1952;38(6):399-421. https://doi.org/10.1016/S0002-9416(52)90128-0
- [10] Lee SJ, Baek SH, Kim SC, Kook YA. Size and forms of the mandibular dental arch in Korean malocclusion patients. Korean J Orthod 2005;35(1):15-22.

- [11] Lee SM, Yoon YJ, Kim KW. Arch form & dimensions after orthodontic treatment by premolar extraction. Korean J Orthod 1998;28(5):717-29.
- [12] Arai K, Will LA. Subjective classification and objective analysis of the mandibular dental-arch form of orthodontic patients. Am J Orthod Dentofacial Orthop 2011;139(4):315-21. https://doi.org/10.1016/j.ajodo.2009.12.032
- [13] Hwang JM, Lee CS, Han JH. The size, form of dental arch and occlusion in dental hygiene students in Korea. J Dent Hyg Sci 2014;14(3):390-6. https://doi.org/10.17135/ jdhs.2014.14.3.390
- [14] Jang JS, Suk KE, Bayome M, Kim TJ, Kim SH, Kook YA. Comparison of arch form between Koreans and Egyptians. Korean J Orthod 2010;40(5):334-41.
- [15] Nam HJ, Son WS, Park SB, Kim SS. Changes of mandibular dental arch during surgicalorthodontic treatment in skeletal class III malocclusion individuals. Korean J Orthod 2008;38(4):283-98.
- [16] Jeon JY, Kim SJ, Kang SG, Park TG. Change in arch width in extraction vs nonextraction treatment. Korean J Orthod 2007;37(1):66-72.
- [17] Lee YH, Yang YM, Lee YH, Kim SH, Kim JG, Baik BJ. A study on the relations of various parts of the palate for primary and permanent dentition. J Korean Acad Pediatr Dent 2004;31(4):569-78.
- [18] Sohn BW, Baik HS. Growth and development of arch form. Korean J Orthod 1998;28(1):17-27.
- [19] Han JH, Lee CS, Hwang JM. Relationship of physique to the size of dental arch in dental hygiene students. J Dent Hyg Sci 2013;13(2):197-202.
- [20] National health insurance service. National health screening statistical yearbook. National health insurance service; [cited 2019 Dec 04]. Available from: http:// www.nhis.or.kr/bbs7/boards/B0159/27944?boardKey=35&sort=sequence&order =desc&rows=10&messageCategoryKey=&pageNumber=1&viewType=generic&ta rgetType=12&targetKey=35&status=&period=&startdt=&enddt=&queryField=&query=
- [21] Hwang JM, Lee CS, Han JH. The size, form of dental arch and occlusion in dental hygiene students in Korea. J Dent Hyg Sci 2014;14(3):390-6.
- [22] Ferrario VF, Sforza C, Miani A Jr, Tartaglia G. Mathematical definition of the shape of dental arches in human permanent healthy dentitions. Eur J Orthod 1994;16(4):287-94. https://doi.org/10.1093/ejo/16.4.287
- [23] Collins BP, Harris EF. Arch form in American blacks and whites with malocclusions. J Tenn Dent Assoc 1998;78(1):15-8.
- [24] Dekock WH. Dental arch depth and width studied longitudinally from 12 years of age to adulthood. Am J Orthod 1992;62(1):56-66. https://doi.org/10.1016/0002-9416(72)90125-x
- [25] Jang KS, Suk KE, Bayome M, Kim YJ, Kim SH, Kook YA. Comparison of arch form between Koreans and Egyptians. Korean J of Orthod 2010;40(5):334-41. https:// doi.org/10.4041/kjod.2010.40.5.334
- [26] Haralabakis NB, Sifakakis I, Papagrigorakis M, Papadakis G. The correlation of sexual dimorphism in tooth size and arch form. World J Orthod 2006;7(3):254-60.
- [27] Harper C. A comparison of medieval and modern dentitions. Eur J Orthod 1994;16(3):163-73. https://doi.org/10.1093/ejo/16.3.163
- [28] Luther F. A cephalometric comparison of medieval skulls with a modern population. Eur J Orthod 1993;15(4):315-25. https://doi.org/10.1093/ejo/15.4.315