Journal of Korean Society of Dental Hygiene

Original Article

Improvement in oral function after an oral exercise program including whole-body exercises

Corresponding Author: Eun-Seo Jung, Department of Dental Hygiene, Shinhan University, 95 Hoam-ro, Uijeongbu, Gyeonggi-do, 11644, Korea. Tel: +82-31-870-3450, +82-10-5415-7913, Fax: +82-31-870-3459, E-mail:

Su-Yeon Seo¹ · Yoon-Young Choi² · Kyeong-Hee Lee³ · Eun-Seo Jung³ ¹Department of Beauty Health Science, Shinhan University ²Artificial Intelligence Big Data Medical Center, Yonsei University Wonju College ³Department of Dental Hygiene, Shinhan University



Received: November 23, 2020 Revised: December 16, 2020 Accepted: December 25, 2020

ABSTRACT

dentalmien@hanmail.net

Objectives: To evaluate the impact of an oral exercise program including whole-body exercises on oral function in older people. **Methods:** The participants (aged \geq 65 years) were divided into three groups: intervention group I (only oral exercise), intervention group II (oral exercise with whole-body exercises), and control group (no intervention). The oral health status, saliva flow rate, and oral muscle strength were evaluated. Analyses were performed to compare the three groups and identify the changes in the aforementioned parameters before and after the program. **Results:** The saliva flow rate significantly increased in intervention group I; the posterior tongue strength and cheek strength also increased but not significantly. The anterior tongue, posterior tongue, and cheek strengths significantly increased in intervention group II. **Conclusions:** The oral exercise program including whole-body exercises showed positive effects on the saliva flow rate and oral exercise strength increased in intervention group II. **Conclusions:** The oral exercise program including whole-body exercises showed positive effects on the saliva flow rate and oral strength. No significant differences were observed in the quality of life related to oral health.

Key Words: Older adults, Oral exercise, Oral function, Whole-body exercises

Introduction

The oral health of older people is one of the very basic factors required for maintaining a healthy life. Oral functions, including chewing, swallowing, and pronunciation, are essential for maintaining social interactions as well as sound quality of life [1]. In particular, oral infection prevention and oral function improvements are necessary in order to masticate

food, enjoy flavours, and display diverse expressions [2]. However, the mouths of elderly individuals often show substantial tooth loss due to oral infections that have accumulated throughout their lives [3]. Furthermore, remaining teeth are often worn out on the occlusal surface due to dental attrition, which results in lower cusps and decreased mastication [4]. In addition, decreased tongue strength, mobility and decreased saliva secretion due to salivary gland aging make swallowing difficult to the elderly [5]. With age-related decreased swallowing, the risk of aspiration (food or saliva entering the airway instead of the oesophagus) increases [6]. If these difficultes persist for a long time, the risk of aspiration pneumonia also increases [7]. Therefore, active efforts are required to improve the factors that negatively affect the quality of life for older people related to their oral health. These include conditions such as decreased chewing, dry mouth, and decreased swallowing [6,7].

The methods for improving the oral health of older people include oral health education, preventative assessments, and early treatment [8]. However, oral health education, which provides the basis for oral health motivation, is less effective due to fewer individuals carrying out the required behavioural changes. Care by experts, preventative assessments, and early treatment also is less effective because they require visits to the oral health centre [8,9]. Therefore, an oral exercise program that can be easily followed in daily life will effectively improve the oral health of older people [10].

In Japan, which has a high life expectancy, oral health improvement methods were added to oral health-related programs through oral health education [11]. Japan has an oral exercise policy for improving oral health [9] that has been reported to be very effective in improving the oral health of older people in an aging and super-aging society [9,11,12]. Furthermore, the recommended exercises can be continuously used by older people without complications, financial burden, or limitations of time or place [13].

An oral exercise program comprised of saliva-producing, chewing-strengthening, swallowingstrengthening, and speaking-strengthening exercises was also developed and applied in older people in facilities throughout Korea. As a result, this program has improved the level of oral health in elderly individuals with healthy oral functions [14]. The program promoted the active recovery, maintenance, and improvement of oral functions in elderly individuals with decreased oral functions. Altogether, it increased the quality of life and oral health of older people [15].

Furthermore, it has been reported that whole-body exercises have aging prevention and health improvement effects in older people [16,17]. These positive effects of whole-body exercises in older people include physiological and affective elements related to decreased depression and anxiety [18] as well as social interactions such as the maintenance of bonding with others. In this respect, whole-body exercises can be considered an essential activity in old age [19]. About 40% of people aged 60 years or older experience discomfort from dysphagia, and this is related to general muscle weakening [20]. Furthermore, whole-body muscle weakening affects muscles of the head and neck as well, and weakening of the tongue muscles is the root cause of dysphagia [21,22]. Therefore, whole-body exercises may have a positive effect on oral health improvements in older people. It is anticipated that oral function improvements will be greater in elderly individuals who perform oral exercises with whole-body exercises than in older adults who only perform oral exercises.

Therefore, an oral exercise program was selected as the oral function improvement methodology of this study. To differentiate our investigation from previous studies, we studied oral function improvements that result from oral exercise combined with whole-body exercise.

Methods

1. Research design

This study applied the oral exercise program with whole body gymnastics to improve oral function to elderly subjects over 65 years of age, and used a non-equivalent pre-post similar experimental design to confirm the effect.

2. Participants

This study was designed as a case-control, three-arm randomized clinical trial. Participants were elderly individuals (aged \geq 65 years) from three centres for senior citizens in Gyeonggi-do, Korea. Individuals with a history of temporomandibular joint dysfunction, oral and maxillofacial surgery, diagnosis of mastication and neck movement limitations, myocardial infarction, and stroke were excluded.

Participant recruitment began on May 1, 2019 and lasted until Dec 31, 2019. The participants were randomly assigned to intervention group I (only oral exercise), intervention group II (oral exercise with whole-body exercise), and the control group (no intervention), by drawing consecutively numbered envelopes.

The required sample size was calculated as 92 individuals using G*power 3.1 program with effect size (d) of 0.25, significance level (α) of 0.05, and test of power (1- β) 85%. The effect size was estimated in reference to a previous study *t*hat used an effect size of 0.25 [23]. Considering drop-out, 40 participants were selected for each group, with a total of 120 participants. After excluding those who terminated their participation during the study, data from a total of 88 participants (32 participants in the intervention group I, 30 in intervention group II, and 26 in the control group) were analyzed <Fig. 1>.

This study was conducted after obtaining approval from 00 University's Institutional Review Board (SHIRB-201902-HR-087-02-(2)). Written consent forms were obtained from the research participants prior to their participation in the study. The study was conducted in full accordance with ethical principles, including the World Medical Association Declaration of Helsinki (version 2008), and the clinical trial was registered with the Clinical Research Information Service of the Korean government (KCT0004827–08/03/2020).



Fig. 1. Consolidated standards of reporting trials(CONSORT) flow diagram

3. Intervention

Detailed information regarding the whole-body exercise and oral exercise program is listed in <Table 1>. The whole-body exercises were performed by controlling breathing and muscle strength in synchrony with a song with a 4/4 rhythm. Oral exercises were applied by referencing the elderly oral function improvement exercises that were developed by a health clinic in Kochi city, Japan [15] and the Korean program [14] For the intervention group I, the oral exercise program was conducted twice a week for five weeks. The participants in the intervention group II performed the oral exercise program with the whole-body exercises twice a week for five weeks.

4. Outcome measures

The general characteristics of the study subjects were sex, age, educational level, economic status, and spouse living together. As for the characteristics of oral health-related behavior, the Frequency of tooth-brushing, tongue cleaning, gum cleaning, use of oral hygiene products, and Scaling during the previous year.

The Oral Health Impact Profile-14 (OHIP-14) [24] was collected through self-reported surveys. OHIP-14 is comprised of a total of 14 questions related to the condition of oral health in the past year. Each item is answered on a 5-point Likert scale. The maximum total score is 70 points, with higher scores indicating better oral health related quality of life. In addition, the Geriatric Oral Health Assessment Index (GOHAI); [25] which is comprised of 12 items, was used to investigate the oral health status of older people over the past three months, with higher scores indicating better oral health related quality of life.

Time	Oral exercise	Time
5 min	I. Oral exercise that promotes saliva production	5 min
	• Move the tongue outside the mouth with the mouth open	
	• Move the tongue inside the mouth with the mouth closed	
5 min	II. Oral exercise to strengthen mastication	5 min
	• Puff up the cheeks as much as possible, then exhale as much air as possible	
	 Stretch the lips, tongue, and cheeks 	
5 min	III. Oral exercise to strengthen swallowing	5 min
	• Breathe in with the nose, stop breathing, and swallow	
	 Breathe out slowly through the mouth 	
5 min	IV. Oral exercise to strengthen talking	5 min
	• Loudly and accurately pronounce: Pa-pa-pa, ta-ta-ta, ka-ka-ka, la-la-la, pa-ta-ka-la	
	Time 5 min 5 min 5 min 5 min	 Time Oral exercise 5 min I. Oral exercise that promotes saliva production Move the tongue outside the mouth with the mouth open Move the tongue inside the mouth with the mouth closed 5 min II. Oral exercise to strengthen mastication Puff up the cheeks as much as possible Stretch the lips, tongue, and cheeks 5 min III. Oral exercise to strengthen swallowing Breathe in with the nose, stop breathing, and swallow Breathe out slowly through the mouth 5 min IV. Oral exercise to strengthen talking Loudly and accurately pronounce: Pa-pa-pa, ta-ta-ta, ka-ka-ka, la-la-la, pa-ta-ka-la

Table 1. Whole-body exercise and oral exercise programs

To measure the unstimulated saliva flow rate, the participants were advised not to take actions that could promote oral functions (talking, eating, drinking, smoking, or oral hygiene) for two hours before collecting saliva. The participants swallowed all the saliva that was in their mouths immediately before the saliva was collected. Then, their saliva was collected in a cup for five minutes. Because this study aimed to identify the changes in the amount of saliva with respect to each exercise program rather than diagnose xerostomia, the amount was measured in weight (g) rather than volume (mL). For this, an ultra-precision electronic scale (Model: BJ-420, Manufacturer: SHINKO) was used. As unstimulated saliva flow rates can differ depending on an individual's overall health status each day, measurements were taken twice, at identical times and places, and the mean value was calculated.

The Iowa Oral Performance Instrument (IOPI)(IOPI Medical, WA, USA) is a portable device for measuring the pressure of anatomical structures in the mouth [26]. This device was used to measure tongue and cheek strength three times, consecutively, and the maximum value (KPa) was recorded. While taking the measurements, a one-minute rest was provided between each exercise to alleviate orofacial muscle fatigue, and all measurements were made by one researcher.

To measure anterior tongue strength, the examiner fixed the bulb so that the alveolar ridge of the hard palate and the region that was 10 mm from the beginning of the tongue were in contact. To measure posterior tongue strength, the examiner fixed the bulb so that the back part of the hard palate and the tongue touched. The participants were subsequently asked to press the bulb toward the hard palate with their tongue, as forcibly as possible, for two seconds. To measure cheek strength, the bulb was placed between the teeth and cheeks, and the participants were asked to softly close their mouth. After placing the bulb in the anterior region of their cheeks, the participants were asked to apply pressure to the bulb, with as much strength as possible, for two seconds.

5. Data analyses

The data collected in this study were analyzed using SPSS Statistics (version 22.0, IBM Corporation, New York, NY, USA), with a statistical significance level of 0.05. To compare the general characteristics and oral health-related behaviours of the research participants, a chi-square test was performed. The ANOVA test was performed to compare the average values after the program of the three groups. To identify the changes that occurred after the program's completion among the three groups, a paired t-test was performed.

Results

1. General characteristics

Sex, age, education level, economic status, and spousal cohabitation were recorded as general characteristics of the research participants. There were no statistically significant differences in these factors among the three groups, showing homogeneity between intervention group I, intervention group II, and the control group <Table 2>. In all three groups, the proportion of female participants was higher than that of male participants, and the 75–84-year age groups contained the greatest participant populations.

Characteristics	Division	Int I	Int II	Con	χ^2	p^{*}
		(n=32)	(n=30)	(n=26)	λ.	1
Sex	Male	4(12.5)	3(10.0)	3(11.5)	0.097	0.953
	Female	28(87.5)	27(90.0)	23(88.5)		
Age (yrs)	65-74	10(31.3)	4(13.3)	7(26.9)	3.003	0.557
	75–84	20(62.5)	23(76.7)	17(65.4)		
	\geq 85	2(6.2)	3(10.0)	2(7.7)		
Education level	No school	2(6.3)	3(10.0)	5(19.2)	4.529	0.339
	Graduated elementary school	11(34.4)	14(46.7)	7(26.9)		
	Graduated middle school or higher	19(59.3)	13(43.3)	14(53.9)		
Economic status	High	2(6.3)	1(3.3)	5(19.2)	7.444	0.114
	Middle	16(50.0)	18(60.0)	8(30.8)		
	Low	14(43.7)	11(36.7)	13(50.0)		
Spouse	Widowed	17(53.1)	12(40.0)	16(61.5)	4.294	0.368
	Cohabitating	14(43.8)	14(46.7)	8(30.8)		
	Separated or divorced	1(3.1)	4(13.3)	2(7.7)		

Table 2. Divisions and contents of the questions

*by chi-square test

Int I; Intervention group I, Int II; Intervention group II, Con; Control group

2. Oral health-related behaviors

The frequency of tooth-brushing, tongue cleaning, gum cleaning, use of oral hygiene products, and scaling in the past year were investigated as factors related to oral health-related behaviours of the research participants. There were no statistically significant differences among the three groups, thereby demonstrating homogeneity among the three groups <Table 3>.

Characteristics	Division	Int I (n=32)	Int II (n=30)	Con (n=26)	χ^2	p^{*}
Frequency of tooth-brushing	1	4(12.5)	6(20.0)	6(23.1)	1.228	0.873
	2	12(37.5)	10(33.3)	9(34.6)		
	≥ 3	16(50.0)	14(46.7)	11(42.3)		
Tongue cleaning	Yes	19(59.4)	18(60.0)	12(46.2)	1.360	0.507
	No	13(40.6)	12(40.0)	14(53.8)		
Gum cleaning	Yes	18(56.3)	21(70.0)	13(50.0)	2.473	0.290
	No	14(43.8)	9(30.0)	13(50.0)		
Usage of oral hygiene product	Yes	15(46.9)	12(40.0)	12(46.2)	0.347	0.841
	No	17(53.1)	18(60.0)	14(53.8)		
Scaling during the previous year	Yes	7(21.9)	7(23.3)	7(26.9)	0.208	0.901
	No	25(78.1)	23(76.7)	19(73.1)		

Table 3. Oral health-related behaviors

*by chi-square test

Int I; Intervention group I, Int II; Intervention group II, Con; Control group

3. Effects of the oral exercise program with whole-body exercise

The findings of the oral function improvement investigations across the two measurements, before and after the program, are as follows <Table 4>. OHIP-14 and GOHAI increased in both intervention groups; however, these increases were not statistically significant. The between-group comparisons of the post-intervention measures also did not show a difference.

Saliva flow rate increased by 0.81 in intervention group I and by 0.42 in intervention group II, and these changes were statistically significant. Furthermore, the between-group comparisons also showed that intervention group I had a greater increase than the other groups (p<0.01).

Anterior tongue strength measured with IOPI increased by 3.28 in intervention group I and by 8.36 in intervention group II, and these changes were statistically significant. Additionally, the between-group comparisons also showed that intervention group II had a greater increase than the other groups (p<0.05). Posterior tongue strength measured with the IOPI increased by 1.93 in intervention group I, however this increase was not statistically significant. Posterior tongue strength also increased by 8.03 in intervention group II, with statistical significance. Furthermore, between-group comparisons also showed that intervention group II had a greater increase in posterior tongue strength than did the other groups (p<0.05). Cheek strength, as measured with the IOPI, increased by 3.09 in intervention group I, without statistical significance, and by 9.00 in intervention group II, with statistical significance. Furthermore, between-group comparisons showed that the increase in cheek strength in intervention group II was greater than that in the other groups (p<0.01). The control group showed changes that were not statistically significant in the various measurements.

Unit: N(%)

Variables	Groups	Pre-intervention	Post-intervention	p^{*}	p^{**}
OHIP-14	Int I	28.21±7.55	31.37±11.19	0.170	0.083
	Int II	30.10±12.93	31.46±10.43	0.497	
	Con	50.96±9.77	46.46±6.58	0.070	
GOHAI	Int I	28.31 ± 8.88	30.40±8.97	0.331	0.158
	Int II	29.63±11.10	30.83±9.90	0.542	
	Con	42.11±8.44	38.84±5.77	0.050	
Saliva flow rate (g)	Int I	1.12 ± 0.68	1.94 ± 1.13	< 0.001	0.002
	Int II	1.11 ± 0.80	1.53 ± 0.93	0.016	
	Con	$0.98 {\pm} 0.80$	$0.98 {\pm} 0.90$	0.919	
Anterior tongue strength (kPa)	Int I	31.18 ± 13.06	34.46±14.72	0.045	0.026
	Int II	25.60 ± 15.97	33.96±16.96	0.001	
	Con	39.50±12.87	38.57±13.17	0.512	
Posterior tongue strength (kPa)	Int I	29.78±14.34	31.72±17.39	0.283	0.012
	Int II	22.40±15.06	30.43±16.77	0.003	
	Con	37.73±14.01	39.65±13.15	0.149	
Cheek strength (kPa)	Int I	20.94±11.86	24.03±13.52	0.266	0.003
-	Int II	23.10±11.78	32.10±19.99	0.001	
	Con	28.65±10.33	26.50±11.16	0.363	

Table 4. Effects of the oral exercise program with whole-body exercise

Unit:Mean±SD

*by paired t-test, **by ANOVA

Int I; Intervention group I, Int II; Intervention group II, Con; Control group

Discussion

This study developed a program that combined whole-body exercises with existing oral exercises and identified related oral function improvements in elderly individuals. Scores of the OHIP-14 and GOHAI evaluations, which are the two representative indicators for measuring the relationship between oral health and quality of life, increased for the intervention groups, without statistical significance. This may be because the program was only five weeks in duration, which could have been too brief a period for individuals to consider that their quality of life improved as a result of their oral health. However, the participants' behavioural changes could become significant when the program is continuously followed. Thus, further long-term measurements and evaluations are required in the future.

Saliva is responsible for keeping the mouth healthy by providing moisture to the oral mucous membranes and to prevent drying [5]. In the mouths of older people, the oral mucous membranes contract and saliva secretion decreases, resulting in xerostomia. Pain in the oral mucous membrane can decrease oral functions, such as mastication, deglutition, and pronunciation, and increase tongue coating, which leads to halitosis [6,7]. The findings of this study showed that the saliva flow rate did not change significantly in the control group, while the individuals in the intervention groups presented significant increases. In particular, the saliva flow rate of intervention group I increased from 1.12 to 1.94, and this change was more pronounced than those of the other groups. Between-group comparisons also showed statistically significant differences. It has been reported that oral

exercises are effective at improving xerostomia and increasing unstimulated saliva flow rates [14]. If appropriate, oral function exercises that are performed before further health deterioration and the functional decline in the salivary glands that accompanies aging and drug complications, can be modified [5]. In conclusion, increased saliva flow rates were found to result only from oral exercises, and it is thought that continuous oral exercise training among older people will have a positive impact on oral functional improvements.

The tongue is a stomatognathic structure that is comprised of various intrinsic and extrinsic muscles [27]. Extrinsic muscles play a role in changing the location of the tongue through protrusion, retraction, depression, and elevation, while intrinsic muscles play a role in changing the shape of the tongue [28]. These muscles cooperate during the deglutition process by regulating the mastication, formation, and manipulation of food in the mouth, and they are also involved in breathing and language [29]. Furthermore, cheek muscles reside in the lower interior location, and the buccinator muscle is the agonist muscle that plays a role in ensuring that food is not left between the teeth and gums (i.e. sulcus lateralis) through continuous contraction during mastication [26]. In this study, the tongue and cheek strength measured with the IOPI showed that anterior tongue strength increased significantly and posterior tongue strength and cheek strength increased without statistically significant differences in intervention group I. However, in intervention group II, anterior tongue strength, posterior tongue strength, and cheek strength significantly increased, and the betweengroup comparisons showed statistically significant differences. In relation to this, previous studies have also implied that oral exercise can improve oral strength, including mastication, the buccinator muscle, and the root of the tongue, due to movement of the oral muscles [26]. Prior studies have also showed that long-term participation in oral exercises with physical activities can inhibit the deterioration of oral functions in older people and contribute to the maintenance of oral health [11]. Tongue and cheek strength were found to significantly improve in this study as well as being partially in line with the findings of previous studies [26]. Furthermore, although it was difficult to clearly hypothesize the reason for the more significant increase in oral strength when whole-body exercises were performed with oral exercises, it may be that relaxation of the body and tense muscles through whole-body exercises can increase blood flow to the brain [30] and increase the effects of oral exercise. Therefore, it is necessary to identify the cause for the observed improvements through future studies.

The findings of this study showed that the saliva flow rate increases with oral exercise alone and that improvements in the saliva flow rate and oral strength were observed when whole-body exercises and oral exercises were performed in parallel. However, because the duration of the study was five weeks, there was not enough time to test the effects of oral exercise with whole-body exercises, and this likely contributed to the lack of large differences between the two groups. Previous research has implied that whole-body exercise with oral exercise contributes to oral function improvements [11]. Considering these findings, it is anticipated that the long-term application of an oral exercise program with whole-body exercise will lead to large oral function improvements.

Although there have been many studies focused on oral exercises as methods for preventing xerostomia in older people, no investigations have used an oral exercise program in combination

with whole-body exercise to improve oral functions. Furthermore, no studies have measured the oral strength using IOPI. Thus, there are difficulties with directly comparing the findings of this study to those of prior studies. This study is significant in that it pioneered an investigation of oral functional improvements for older adults with insufficient prior research. This study also compared the differences in the oral functional improvements of regular oral exercise with oral exercise in combination with whole-body exercise. It is thought that an oral exercise program with whole-body exercise will increase the oral health related quality of life of older people and contribute to oral health improvements. This combined program will benefit an aging society.

Conclusions

To evaluate the impact of an oral exercise program with whole-body exercise on the oral function of older people. The participants (aged \geq 65 years) were divided into three groups: intervention group I (only oral exercise), intervention group II (oral exercise with whole-body exercise), and control group. Before and after the program, the oral health status, saliva flow rate, and oral muscle strength were measured. Analyses were performed to compare the three groups and to identify changes in the three groups before and after the program.

1. Saliva flow rate increased in intervention group I and in intervention group II. These increases were statistically significant.

2. In oral strength measurements using the IOPI, anterior tongue strength increased significantly in intervention group I. In this group, posterior tongue and cheek strength increased; however, these differences were not statistically significant.

3. In intervention group II, anterior tongue, posterior tongue, and cheek strength increased, and these changes were statistically significant.

The oral exercise program with whole-body exercise was found to have positive effects on increasing the saliva flow rate and oral strength. No significant differences were observed in the quality of life related to oral health. Therefore, in subsequent studies, repeated studies are needed to confirm changes in oral health quality of life through long-term measurement.

Acknowledgements

This study was funded by the National Research Foundation of Korea and supported by the 2019 Ministry of Science and Information Communication Technology (No. 2018R1A2B6006701). The funders had no role in the study design, data collection and analysis, or preparation of the manuscript.

Conflicts of Interest

The authors declared no conflict of interest.

Authorship

Conceptualization: KH Lee, ES Jung, YY Choi, SY Seo; Data collection: KH Lee, ES Jung, SY Seo; Formal analysis: ES Jung; Writing - original draft: ES Jung; Writing - review & editing: YY Choi, SY Seo

References

- Lee SH, Shin BM, Shin SJ, Bae SM. Relationship between perceived health status and health-related quality of life in the elderly: a focus on moderating effects of oral health. J Korean Soc Dent Hyg 2019;19(5):715-29. https://doi.org/10.13065/jksdh.20190061
- [2] Moon SH, Hong GR. Predictors of chewing discomfort among community-dwelling elderly. J Korean Acad Community Health Nurs 2017;28(3):302-12. https://doi.org/10.12799/jkachn.2017.28.3.302
- [3] Yang SO, Jeong GH, Kim SJ, Kim KW, Lee SH, Saung SY, et al. The effects of oral function improving exercise on the UWS, oral function and OHIP in elderly. J Korean Public Health Nurs 2012;26(3):478-90. https://doi.org/10.5932/JKPHN.2012.26.3.478
- [4] Gérald M, Saliba-Serre B, Ferrandez AM, Ruquet M. Objective and perceived oral health status of elderly nursing home residents: a local survey in southern France. Clin Interv Aging 2019;28(14):1141-51. https://doi.org/10.5932/10.2147/CIA.S204533
- [5] Kim JH, Kim HY. Effects of an oral self-care program on the elderly's xerostomia and oral health-related quality of life. J Korean Acad Community Health Nurs 2018;29(3):382-92. https:// doi.org/10.12799/jkachn,2018.29.3.382
- [6] Dalodom S, Lam-Ubol A, Jeanmaneechotechai S, Takamfoo L, Intachai W, Duangchada K, et al. Influence of oral moisturizing jelly as a saliva substitute for the relief of xerostomia in elderly patients with hypertension and diabetes mellitus. Geriatr Nurs 2016;37(2):101-9. https:// doi.org/10.1016/j.gerinurse.2015.10.014
- [7] Tanigawa T, Yamashita J, Sato T, Shinohara A, Shibata R, Ueda H, et al. Efficacy and safety of pilocarpine mouthwash in elderly patients with xerostomia. Spec Care Dentist 2015;35(4):164-9. https://doi.org/10.1111/scd.12105
- [8] Lee SH, Ryu JA, Yu HE, Lee JH, Shin SJ. Comparison of effects according to type of oral exercise program for elderly in Gangneung city. J Dent Hyg Sci 2016;16(6):424-31. https://doi.org/10.17135/ jdhs.2016.16.6.424
- [9] Ibayashi H, Fujino Y, Pham TM, Matsuda S. Intervention study of exercise program for oral function in healthy elderly people. Tohoku J Exp Med 2008;215(3):237-45. https://doi.org/10.1620/tjem.215.237
- [10] Jang KA, Hwang IC. Objective effects and satisfaction of mouth gymnastics program. J Korea Cont Assoc 2011;11(10):388-95. https://doi.org/10.5392/JKCA.2011.11.10.388
- [11] Miyoshi S, Shigeishi H, Fukada E, Nosou M, Amano H, Sugiyama M. Association of oral function with long-term participation in community-based oral exercise programs in older Japanese women: A cross-sectional study. J Clin Med Res 2019;11(3):165-70. https://doi.org/10.14740/jocmr3664
- [12] Sugiyama T, Ohkubo M, Honda Y, Tasaka A, Nagasawa K, Ishida R, et al. Effect of swallowing exercises in independent elderly. Bull Tokyo Dent Coll 2013;54(2):109-15. https://doi.org/10.2209/ tdcpublication.54.109
- [13] Jeon YJ, Choi JS, Han SJ. The effect of dry mouth improvement by oral exercise program in elderly people. J Korean Soc Dent Hyg 2012;12(2):293-305. https://doi.org/10.13065/jksdh.2012.12.2.293
- [14] Hwang YS, Cho EP. A study on elderly people's satisfaction level with oral exercise program. J Korean Soc Dent Hyg 2009;9(4):795-807.

- [15] Hirosaki M, Ohira T, Kajiura M, Kiyama M, Kitamura A, Sato S, et al. Effects of a laughter and exercise program on physiological and psychological health among community-dwelling elderly in Japan: Randomized controlled trial. Geriatr Gerontol Int 2013;13(1):152-60. https://doi.org/10.1111/j.1447-0594.2012.00877.x
- [16] Lee KY, Yoo KS, Kim HO, Yang YR. The effects of shinbaram health exercise on health status and daily living performance ability of the elderly. Korean Pub Health Res 2018;44(3):89-103. https:// doi.org/10.22900/kphr.2018.44.3.007
- [17] Kim SH, An EJ. A study on developing gymnastic program for seniors with mild dementia & walking problems : measuring its effects. J Sport Leis Stud 2015;59:601-18.
- [18] Kim JE. An analysis on the structural relationship among subjective value, sentiment and psychological well-being of the elderly participating in the gymnastics for all. J Korean Soc Sports Sci 2014;23(1):501-12.
- [19] Hong MH. The relationship of self expression, self efficacy and psychological happiness of the elderly participating in daily exercise. J Korean Phys Edu Assoc Girls Women 2016;30(4):183-97. https:// doi.org/10.16915/jkapesgw.2016.12.30.4.183
- [20] Doggett DL, Turkelson CM, Coates V. Recent developments in diagnosis and intervention for aspiration and dysphagia in stroke and other neuromuscular disorders. Curr Atheroscler Rep 2002;4:311-8. https://doi.org/10.1007/s11883-002-0010-x
- [21] Robbins JA, Levine R, Wood J, Roecker EB, Luschei E. Age effects on lingual pressure generation as a risk factor for dysphagia. J Gerontol A Biol Sci Med Sci 1995;50(5):257-62. https://doi.org/10.1093/ gerona/50a.5.m257
- [22] Nicosia MA, Hind JA, Roecker EB, Carnes M, Doyle J, Dengel GA, et al. Age effects on the temporal evolution of isometric and swallowing pressure. J Gerontol A Biol Sci Med Sci 2000;55(11):634-40. https://doi.org/10.1093/gerona/55.11.m634
- [23] Lee KY. Development and evaluation of professional oral care program for elderly in long-term care facilities[Doctoral dissertation]. Cheonan: Univ. of Namseoul, 2016.
- [24] Locker D, Matear D, Stephens M, Lawrence H, Payne B. Comparison of the GOHAI and OHIP-14 as measures of the oral health-related quality of life of the elderly. Community Dent Oral Epidemiol 2001;29(5):373-81. https://doi.org/10.1034/j.1600-0528.2001.290507.x
- [25] Atchison KA, Dolan TA. Development of the geriatric oral health assessment index. J Dent Educ 1990;54(11):680-7. https://doi.org/10.1002/j.0022-0337.1990.54.11.tb02481.x
- [26] Adams V, Mathisen B, Baines S, Lazarus C, Callister R. Reliability of measurements of tongue and hand strength and endurance using the iowa oral performance instrument with healthy adults. Dysphagia 2014;29(1):83-95. https://doi.org/10.1007/s00455-013-9486-5
- [27] Yano J, Yamamoto-Shimizu S, Yokoyama T, Kumakura I, Hanayama K, Tsubahara A. Effects of anterior tongue strengthening exercises on posterior tongue strength in healthy young adults. Arch Oral Biol 2019;98:238-42. https://doi.org/10.1016/j.archoralbio.2018.11.028
- [28] Trawitzki LVV, Borges CGP, Giglio LD, Silva JB. Tongue strength of healthy young adults. J Oral Rehabil 2011; 38(7):482-6. https://doi.org/10.1111/j.1365-2842.2010.02182.x
- [29] Kim CY, Lee YM, Ha EH. Development and evaluation of a dysphagia assessment tool and an intervention program for the elderly in the long-term care facilities. J Korea Acad-Indus Cooper Soc 2012;13(2):685-93. https://doi.org/10.5762/KAIS.2012.13.2.685
- [30] Weijenberg RAF, Scherder EJA, Lobbezoo F. Mastication for the mind-the relationship between mastication and cognition in ageing and dementia. Neurosci Biobehav Rev 2011;35(3):483-97. https://doi.org/10.1016/j.neubiorev.2010.06.002